

FERMION SYSTEMS WITH FUZZY SYMMETRIES

(LEVERAGING THE KNOWN TO UNDERSTAND THE UNKNOWN)

J. P. DRAAYER¹, K. D. SVIRATCHEVA¹,
T. DYTRYCH¹, C. BAHRI¹, K. DRUMEV¹, J. P. VARY²

¹*Department of Physics and Astronomy, Louisiana State University,
Baton Rouge, Louisiana 70803, USA*

²*Department of Physics and Astronomy, Iowa State University,
Ames, IA 50011, USA*

Microscopic models, which embody the simplicity and significance of a dynamical symmetry approach to nuclear structure, are reviewed. They can reveal striking features of atomic nuclei when a symmetry dominates and solutions in domains that may otherwise be unreachable.

1. Overview of algebraic fermion models. A theory that invokes group symmetries is driven by an expectation that the wave functions of the quantum mechanical system under consideration can be characterized by their invariance properties under the corresponding symmetry transformations. But even when the symmetries are not exact, if one can find near invariant operators, the associated symmetries can be used to help reduce the dimensionality of a model space to a tractable size. Throughout the years, group-theoretical approaches have identified fundamental symmetries in light to heavy nuclei and achieved a reasonable reproduction of experimental data (for a review of fermion models, see ¹). In addition, they provide theoretical predictions for nuclear systems including heavy unstable nuclei not yet explored, and ‘exotic’ nuclei, such as neutron-deficient or $N \approx Z$ nuclei on the path of the nucleosynthesis rp -processes.

It is well-known that effective two-body interactions in nuclei are dominated by pairing and quadrupole terms. The former gives rise to a pairing gap in nuclear spectra, and the latter is responsible for enhanced electric quadrupole transitions in collective rotational bands. Indeed, within the framework of the harmonic oscillator shell-model, both limits have a clear algebraic structure in the sense that the spectra exhibit a dynamical symmetry. In the pairing limit the symplectic $Sp(4)$ ($\sim SO(5)^{2,3}$) group together

with its dual $\text{Sp}(2\Omega)$, for 2Ω shell degeneracy, use the seniority quantum number^{4,5} to classify the spectra. On the other hand, in the quadrupole limit the $\text{SU}(3)$ (sub-) structure⁶ governs a shape-determined dynamics.

In light deformed nuclei, $A \lesssim 28$, the Elliot's $\text{SU}(3)$ model⁶, which incorporates the particle quadrupole and angular momentum operators, proved successful for a microscopic description of collective modes. Indeed, $\text{SU}(3)$ is the exact symmetry group of the spherical oscillator, which is a reasonable approximation for the average potential experienced by nucleons in nuclei. Also, $\text{SU}(3)$ is the dynamical symmetry group of the deformed oscillator, when, as is usually the case, the deformation is generated by quadrupole interactions. In many cases, a single-irrep or few-irreps calculations suffice to achieve good agreement with experimental rotational energy spectra and electromagnetic transitions (e.g., see ^{7,8}).

Limitations due to the fact that the $\text{SU}(3)$ model is applied within a shell and in turn requires effective charges for transition strengths are overcome in the $\text{Sp}(3, \mathbb{R})$ symplectic shell model for light nuclei (for a review see ⁹). It embeds the $\text{SU}(3)$ symmetry and in addition introduces important inter-shell excitations, including high- $\hbar\omega$ correlations and core excitations. The symplectic shell model is a microscopic formulation of the Bohr-Mottelson collective geometric model with a direct relation between a second- and a third-order scalar products of the quadrupole operator and the (β, γ) -shape variables. The $\text{Sp}(3, \mathbb{R})$ symplectic model provides a microscopic description of monopole and quadrupole collective modes in deformed nuclei and a reproduction of experimental rotational energy spectra and electromagnetic transitions without effective charges (e.g., see ^{10,11}).

Furthermore, in the domain of light nuclei one can combine the $\text{Sp}(3, \mathbb{R})$ symplectic shell model and the no-core shell model (NCSM)¹² to push forward the present frontiers in nuclear structure physics. The NCSM+ $\text{Sp}(3, \mathbb{R})$ allows us to use modern realistic interactions without any approximation (for the interaction and the size of the model space) for low- $\hbar\omega$ configurations and hence to fully account for important short- and intermediate-range correlations, while selecting only dominant high- $\hbar\omega$ basis states responsible for multi-shell development of collective motion.

In the region of medium mass nuclei around the $N = Z$ line (currently explored by radioactive beam experiments) protons and neutrons occupy the same major shell and hence their mutual interactions are expected to strongly influence the structure and decay modes of such nuclei. In addition to like-particle (pp and nn) pairing correlations the close interplay of pp , nn and proton-neutron (pn) pairs and the isospin-symmetry influence are

microscopically described by the $\text{Sp}(4)$ pairing model¹³.

For heavy nuclei ($A \gtrsim 100$), the discovery of the pseudo-spin symmetry^{14,15} and its fundamental nature^{16,17,18} establishes the pseudo-SU(3) model¹⁹. The pseudo-spin scheme is an excellent starting point for a many-particle description of heavy nuclei, whether or not they are deformed. As for the SU(3) shell model, in many cases leading-irrep calculations (e.g., see ²⁰) or mixed-irrep calculations (e.g., see ²¹) achieve good agreement with experimental data. The pseudo-SU(3) shell model provides a further understanding of the $M1$ transitions in nuclei such as the even-even ^{160–164}Dy and ^{156–160}Gd isotopes, specifically it reflects on the scissors and twist modes as well as the observed fragmentation, that is, the break-up of the $M1$ strength among several levels closely clustered around a few strong transition peaks in the 2-4 MeV energy region²².

In medium-mass and heavy nuclei, where the pseudo-spin scheme can be applied to the normal parity orbitals and valence spaces are intruded by a unique parity highest- j orbit from the shell above, a major step towards understanding the significance of the intruder level is achieved by a pseudo-SU(3) plus intruder level shell model²³ that is currently under development²⁴.

Furthermore, the advantages of the symplectic $\text{Sp}(3, \mathbb{R})$ extension of the SU(3) model can be employed beyond the light nuclei domain towards a description of heavy nuclei in the framework of the pseudo- $\text{Sp}(3, \mathbb{R})$ shell model (e.g., see ²⁵). While early developments demonstrate the potential of such a model for studying the structure of heavy nuclear systems, it has not been fully explored.

In what follows, we present some recent results for three algebraic fermion models where the symmetries are fuzzy – meaning not exact – but nevertheless extremely useful in gaining a deeper understanding of the structure of real nuclei.

2. Symplectic $\text{Sp}(4)$ pairing model. An algebraic approach, with $\text{Sp}(4)$ the underpinning symmetry and with only six parameters, can be used to provide a reasonable microscopic description of pairing-governed 0^+ states in a total of 306 even-even and odd-odd nuclei with mass $40 \leq A \leq 100$ where protons and neutrons are filling the same major shell^{26,13}. We employ the most general Hamiltonian with $\text{Sp}(4)$ dynamical symmetry,

$$\begin{aligned}
 H_{\text{sp}(4)} = & -G \sum_{i=-1}^1 \hat{A}_i^\dagger \hat{A}_i - F \hat{A}_0^\dagger \hat{A}_0 - \frac{E}{2\Omega} (\hat{T}^2 - \frac{3\hat{N}}{4}) \\
 & - D(\hat{T}_0^2 - \frac{\hat{N}}{4}) - C \frac{\hat{N}(\hat{N}-1)}{2} - \epsilon \hat{N},
 \end{aligned} \tag{1}$$

where G, F, E, D, C and $\epsilon > 0$ are parameters (refer to Table I in ¹³ for their estimates). In (1), $\hat{N}$ counts the total number of particles, $\hat{T}^2$ is the isospin operator and the $\hat{A}_{0,+1,-1}^\dagger$ group generators, which build the basis states by a consequent action on the vacuum state (a core like ^{40}Ca or ^{56}Ni), create, respectively, a proton-neutron (pn) pair, a proton-proton (pp) pair or a neutron-neutron (nn) pair of total angular momentum $J^\pi = 0^+$ and isospin $T = 1$. The model Hamiltonian (1) represents an effective microscopic interaction that conserves the T_0 third projection of the isospin and includes proton-neutron and like-particle isovector ($T = 1$) pairing plus symmetry terms (the latter is related to a proton-neutron isoscalar ($T = 0$) force). The model interaction (1) is found to correlate strongly with realistic interactions like CD-Bonn+3 terms²⁷ in the $1f_{7/2}$ region²⁸ and reflects a large portion of the GXPF1 realistic interaction²⁹ in the upper- fp shell. In addition, the D -term in (1) introduces isospin symmetry breaking and the F -term accounts for a plausible, but weak, isospin mixing³⁰. Both terms are significant in non-analog β -decays studies and also yield quantitative results that are better (e.g., by 85% in $1f_{7/2}$) than the ones with $F = D = 0$.

Good agreement with experiment (small χ -statistics) is observed (Fig. 1) when theoretical eigenvalues of (1) are compared to Coulomb-corrected³¹ experimental energies³². The theory predicts the lowest isobaric analog 0^+ state energy with a deviation ($\chi/\Delta E_{0,exp} \times 100[\%]$) of 0.5% for $1f_{7/2}$ and $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ nuclei in the corresponding energy range considered, $\Delta E_{0,exp}$. The model estimates the binding energy of the proton-rich ^{48}Ni nucleus to be 348.1 MeV, which is 0.07% greater than the sophisticated semi-empirical estimate³³ and only 4% away from the experimental value reported later-on³⁴. The ^{68}Se waiting-point nuclide along the rp -path is estimated to be 574.3 MeV, only 2.7% away from the 2004 precise mass measurement³⁵. Likewise, for the odd-odd nuclei with energy spectra not yet measured the theory predicts the energy of their lowest isobaric analog 0^+ state: 358.62 MeV (^{44}V), 359.34 MeV (^{46}Mn), 357.49 MeV (^{48}Co), 394.20 MeV (^{50}Co) (Fig. 1, right). The $\text{Sp}(4)$ model predicts the relevant 0^+ state energies for an additional 165 even- A nuclei in the medium mass region (Fig. 1, left). The binding energies for 25 of them are also calculated in ³³. For these even-even nuclei, we predict binding energies that on average are 0.05% less than the semi-empirical approximation³³.

Furthermore, without any parameter variation, the theoretical energy spectra of the isobaric analog 0^+ states are found to agree remarkably well with the experimental values where data is available¹³. This agreement represents a valuable result because the higher-lying 0^+ states under




Figure 1. Theoretical isobaric analog 0^+ state energies in MeV of isobaric sequences (lines) (including the Coulomb energy) vs. the isospin projection T_0 . Left: isobars with $A = 56, 58, \dots, 100$ in the $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ shell (^{56}Ni core), compared to experimental values (black 'x') and semi-empirical estimates by P. Möller *et al.* (blue '+'). Right: isobars in the $1f_{7/2}$ level compared to experimental binding energies (x) or energies of the lowest isobaric analog 0^+ excited states (o).

consideration constitute an experimental set independent of the data that enters the statistics to determine the model parameters in (1).

In addition, we examine the detailed features of nuclei by discrete derivatives of the energy function (1) filtering out the strong mean-field influence³⁶. This investigation reveals a remarkable reproduction of the two-proton S_{2p} and two-neutron S_{2n} separation energies, the irregularities found around the $N = Z$ region, the like-particle and pn isovector pairing gaps, and a prominent staggering behavior observed between groups of even-even and odd-odd nuclides³⁶. The zero point of S_{2p} along an isotone sequence determines the two-proton-drip line, which according to the $Sp(4)$ model for the $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ shell lies near the following even-even nuclei³⁶: $^{60}\text{Ge}_{28}$, $^{64}\text{Se}_{30}$, $^{68}\text{Kr}_{32}$, $^{72}\text{Sr}_{34}$, $^{76}\text{Zr}_{36}$, $^{78}\text{Zr}_{38}$, $^{82}\text{Mo}_{40}$, $^{86}\text{Ru}_{42}$, $^{90}\text{Pd}_{44}$, $^{94}\text{Cd}_{46}$, beyond which the higher- Z isotones are unstable with respect to diproton emissions in close agreement with other estimates^{37,33,38} despite the lack of experimental data. In addition, we find a small quadratic mean of the difference in S_{2p} between our model and the other theoretical predictions where data is available, namely, 0.32 MeV in comparison with ³⁷, 0.78 MeV with ³³ and 0.43 MeV with ³⁸.

While the model describes only isobaric analog 0^+ states of even- A medium mass nuclei with protons and neutrons in the same shell, it reveals a fundamental feature of the nuclear interaction, which governs these states. Namely, the latter possesses a clear $Sp(4)$ dynamical symmetry.

3. Pseudo-SU(3) plus intruder level model. The role of intruder levels that penetrate down into lower-lying shells in atomic nuclei has been the focus of many studies and debates. These levels are found in heavy deformed nuclei where the strong spin-orbit interaction destroys an underlying harmonic oscillator symmetry of the nuclear mean-field potential.

We carry out m -scheme shell-model calculations for the ^{58}Cu and ^{64}Ge nuclei in the $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ model space assuming the occupancy of the $f_{7/2}$ orbital to be ‘frozen’. This choice was motivated by the $f_{7/2}$ orbit’s high occupation as reported elsewhere³⁹. The Hamiltonian we used is a G -matrix with a phenomenologically adjusted monopole part⁴⁰. A renormalized




Figure 2. Energy spectrum and B(E2) transition strengths for ^{64}Ge . The width of the arrows in the figure represent the relative B(E2) strengths, normalized to unity for the ground band $2^+ \rightarrow 0^+$ transition. Numbers in each box are for the $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ and the restricted upper fp model space, respectively.

Results for the energy spectrum and B(E2) strengths of ^{64}Ge are shown in Fig. 2 for both model spaces. The renormalized version of the theory in the upper fp subspace not only reproduces the excitation energies obtained in the larger $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ space, but also gives very similar values for the B(E2) transition strengths. These results also confirm those from a study using a schematic interaction⁴². Similar behavior was observed for ^{58}Cu . Besides the ground state and gamma bands for ^{64}Ge a new (possibly beta) band is identified.

In addition, after rescaling occupancies of the states obtained in the upper fp subspace, a pattern that is very similar to that of the occupancies in the larger $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ space is obtained.

In short, novel shell-model calculations for ^{58}Cu and ^{64}Ge in the $1f_{5/2}2p_{1/2}2p_{3/2}1g_{9/2}$ model space using a realistic interaction are compared to those generated by an appropriately renormalized counterpart of the interaction in the truncated upper fp subspace. The results suggest that reliable computations can be performed in a space that does not explicitly include the intruder level as long as the interaction and the transition operators are renormalized appropriately.

4. No-core shell model plus $\text{Sp}(3, \mathbb{R})$ extension. The symplectic shell model is based on the noncompact symplectic $\mathfrak{sp}(3, \mathbb{R})$ algebra with a subalgebraic structure that gives rise to rich underlying physics for a microscopic description of multiple collective modes in nuclei. This follows from the fact that the mass quadrupole and monopole moments operators, the many-particle kinetic energy, the angular and vibrational momenta are all elements of the $\mathfrak{sp}(3, \mathbb{R}) \subset \mathfrak{su}(3)$ algebraic structure⁹.

The symplectic basis states, $|\Gamma_\sigma \Gamma_n \rho \Gamma_\omega \kappa(LS)JM_J\rangle$, are constructed by acting with polynomials in the symplectic raising operator, $A^{(20)}$, on a set of basis states of a symplectic bandhead, $|\Gamma_\sigma\rangle$. They are labeled according to the reduction chain

$$\begin{array}{ccccc} \text{Sp}(3, \mathbb{R}) & \supset & \text{U}(3) & \supset & \text{SO}(3) \\ \Gamma_\sigma & & \Gamma_n \rho & & \Gamma_\omega \quad \kappa \quad L \end{array}$$

where $\Gamma_\sigma \equiv N_\sigma(\lambda_\sigma, \mu_\sigma)$ labels $\text{Sp}(3, \mathbb{R})$ irreducible representations. The $\Gamma_n \equiv n(\lambda_n, \mu_n)$ set of quantum numbers gives the overall $\text{SU}(3)$ coupling of $n/2$ raising operators acting on $|\Gamma_\sigma\rangle$. $\Gamma_\omega \equiv N_\omega(\lambda_\omega, \mu_\omega)$ specifies the $\text{SU}(3)$ symmetry of a symplectic state and $N_\omega = N_\sigma + n$ is the total number of oscillator quanta related to the eigenvalue, $N_\omega \hbar\Omega$, of a center-of-mass motion free harmonic-oscillator (HO) Hamiltonian. The basis states of a $\text{Sp}(3, \mathbb{R})$ irreducible representation can be expanded in HO (m -scheme) basis, which is the basis utilized by the no-core shell-model (NCSM)¹².

In the case of ^{12}C one can construct 13 unique so-called $0\hbar\Omega$ - $\text{Sp}(3, \mathbb{R})$ irreducible representations. $0\hbar\Omega$ means that the symplectic bandhead basis states, $|\Gamma_\sigma\rangle$, lie within $0\hbar\Omega$ many-particle harmonic-oscillator space. For each of the $0\hbar\Omega$ - $\text{Sp}(3, \mathbb{R})$ irreducible representations we generate basis states up to $N_{max} = 6(6\hbar\Omega)$, which is the current limit for NCSM calculations. Typical the dimension of a symplectic representation is of order 10^3 ,

comparing to 10^7 in the case of NCSM m -scheme basis space.

Table 1. Probability distribution of the three most important $\text{Sp}(3, \mathbb{R})$ irreps and of the NCSM wavefunctions.

J=0					
	$0\hbar\Omega$	$2\hbar\Omega$	$4\hbar\Omega$	$6\hbar\Omega$	Total
$(0\ 4)S = 0$	46.26	11.39	4.15	1.11	62.91
$(1\ 2)S = 1$	4.80	1.87	0.80	0.30	7.77
$(1\ 2)S = 1$	4.72	1.84	0.79	0.29	7.64
$\text{Sp}(3, \mathbb{R})$ Total	55.78	15.10	5.74	1.70	78.32
NCSM	56.18	22.40	12.81	7.00	98.38
J=2					
$(0\ 4)S = 0$	46.80	11.33	3.99	1.06	63.18
$(1\ 2)S = 1$	4.84	1.65	0.69	0.25	7.43
$(1\ 2)S = 1$	4.69	1.60	0.67	0.24	7.20
$\text{Sp}(3, \mathbb{R})$ Total	56.33	14.58	5.35	1.55	77.81
NCSM	56.63	21.79	12.73	7.28	98.43
J=4					
$(0\ 4)S = 0$	51.45	11.23	3.71	0.94	67.33
$(1\ 2)S = 1$	3.04	0.89	0.35	0.12	4.40
$(1\ 2)S = 1$	3.01	0.88	0.35	0.12	4.36
$\text{Sp}(3, \mathbb{R})$ Total	57.50	13.00	4.41	1.18	76.09
NCSM	57.64	20.34	12.59	7.66	98.23

The lowest-lying eigenstates of ^{12}C are calculated by NCSM approach using the Many Fermion Dynamics (MFD) code⁴³ with the effective interaction derived from the realistic JISP16 NN potential for oscillator strengths of $\hbar\Omega = 15$ MeV. The large overlaps of the symplectic states with the NCSM wave functions for 0, 2, 4 and $6\hbar\Omega$ subspaces of the m -scheme basis (Tables 1) reveal that around 80% of the latter are symmetric under $\text{Sp}(3, \mathbb{R})$ transformations. Apparently, for all three wave functions, the highest contribution comes from the leading, most deformed, $(0\ 4)S = 0$ $\text{Sp}(3, \mathbb{R})$ irreducible representation. This contribution gets higher towards $J = 4_1^+$, where mixing due to other, less deformed, configurations decreases.

Clearly, the 0^+ , 2^+ and 4^+ states, which are constructed in terms of the three $\text{Sp}(3, \mathbb{R})$ irreps with probability amplitudes defined by the overlaps with the NCSM wavefunctions, can be used as a quite good approximation for a microscopic description of the 0_{gs}^+ , 2_1^+ and 4_1^+ states in ^{12}C . Within this assumption, the $B(E2 : 2_1^+ \rightarrow 0_{gs}^+)$ value turns out to be as much as 81% of the NCSM estimate.

In short, the low-lying states in ^{12}C are quite well explained by only three $\text{Sp}(3, \mathbb{R})$ irreps of 1098 symplectic states, that is only 0.003% of the

NCSM space dimension, with a dominance of the most deformed $(0\ 4)S = 0$ collective configuration. Our findings, as a ‘proof-of-principle’, suggests that a $\text{NCSM} + \text{Sp}(3, \mathbb{R})$ structure could allow one to extend no-core calculations to higher $\hbar\omega$ and heavier nuclei.

In summary, models based on exact or just good (broken but dominant) symmetries in fermion systems can play a significant role in our understanding low-lying nuclear structure; specifically, as shown here, in the development of collective rotational motion and the formation of correlated pairs in nuclei. They also allow us to truncate a model space to typically only a fraction the size encountered in models that do not exploit what we have dubbed here as fuzzy symmetries.

Acknowledgments

This work was supported by the US National Science Foundation, Grant Numbers 0140300 & 0500291 and the Southeastern Universities Research Association (SURA).

References

1. J. P. Draayer, *Fermion Models*, in *Algebraic Approaches to Nuclear Structure*, ed. R. Casten, (Harwood Academic Publishers), Ch. 7, 423 (1992).
2. K. Helmers, *Nucl. Phys.* **23**, 594 (1961).
3. K. T. Hecht, *Nucl. Phys.* **63**, 177(1965); *Phys. Rev.* **139**, B794 (1965); *Nucl. Phys.* **A102**, 11 (1967); J. N. Ginocchio, *Nucl. Phys.* **74**, 321 (1965).
4. G. Racah, *Phys. Rev.* **62**, 438 (1942); *Phys. Rev.* **63**, 367 (1943).
5. B. H. Flowers, *Proc. Roy. Soc. (London)* **A212**, 248 (1952).
6. J. P. Elliott, *Proc. Roy. Soc. (London)* **A245**, 128 (1958); **A245**, 562 (1958); J. P. Elliott and M. Harvey, *Proc. Roy. Soc. (London)* **A272**, 557 (1963).
7. H.A. Naqvi and J.P. Draayer, *Nucl. Phys.* **A516**, 351 (1990).
8. C.E. Vargas, J.G.Hirsch, and J.P.Draayer, *Nucl. Phys.* **A690**, 409 (2001) .
9. D.J. Rowe, *Rep. Prog. Phys.* **48**, 1419 (1985).
10. G. Rosensteel and D.J. Rowe, *Phys. Rev. Lett.* **38**, 10 (1977); *Ann. Phys. N Y* **126**, 343 (1980).
11. J.P. Draayer, K.J. Weeks, and G. Rosensteel, *Nucl. Phys.* **A 413**, 215 (1984).
12. P. Navrátil, J.P. Vary, and B.R. Barrett, *Phys. Rev. Lett.* **84**, 5728 (2000).
13. K. D. Sviratcheva, A. I. Georgieva, and J. P. Draayer, *Phys. Rev.* **C70**, 064302 (2004).
14. K. T. Hecht and A. Adler, *Nucl. Phys.* **A137**, 129 (1969).
15. A. Arima A, M. Harvey, and M K. Shimizu, *Phys. Lett.* **30B**, 517 (1969).
16. J.P. Draayer, in *Proceedings of Symmetries in Physics*, Cocoyoc, Mexico, June 3-7, 1991, ed. A. Frank and K.B. Wolf (Springer-Verlag, Berlin).
17. C. Bahri, J. P. Draayer, and S.A. Moszkowski, *Phys. Rev. Lett.* **68**, 2133 (1992).

18. A.L. Blokhin, C. Bahri, and J.P. Draayer, *Phys. Rev. Lett.* **74**, 4149 (1995).
19. R.D. Ratna Raju, J.R. Draayer, and K.T. Hecht, *Nucl. Phys.* **A202**, 433 (1973).
20. J.R. Draayer and K.J. Weeks, *Phys. Rev. Lett.* **51**, 1422 (1983).
21. G. Popa, J. G. Hirsch, and J. P. Draayer, *Phys. Rev.* **C62**, 064313 (2000).
22. T. Beuschel, J. P. Draayer, D. Rompf, and J. G. Hirsch, *Phys. Rev.* **C57**, 1233 (1998).
23. K.J. Weeks, C.S. Han, and J.P. Draayer, *Nucl. Phys.* **A371**, 19 (1981).
24. K. P. Drumev, C. Bahri, V. G. Gueorguiev and J. P. Draayer, in *Proceedings of Nuclear Physics, Large and Small: International Conference on Microscopic Studies of Collective Phenomena*, Morelos, Mexico, April 19-22, 2004, eds. R. Bijker, R.F. Casten, and A. Frank, Melville, New York, *AIP Conference Proceeding* **726**, 219 (2004).
25. O. Castaños, P.O. Hess, J.P. Draayer, and P. Rochford, *Nucl. Phys.* **A524**, 469 (1991).
26. K. D. Sviratcheva, A. I. Georgieva, and J. P. Draayer, *J. Phys. G: Nucl. Part. Phys.* **29**, 1281 (2003).
27. S. Popescu, S. Stoica, J. P. Vary, and P. Navratil, to be published.
28. K. D. Sviratcheva, J. P. Draayer, and J. P. Vary, *Phys. Rev.* **C73**, 034324 (2006).
29. M. Honma, T. Otsuka, B. A. Brown, and T. Mizusaki, *Phys. Rev.* **C69**, 034335 (2004).
30. K. D. Sviratcheva, A. I. Georgieva, and J. P. Draayer, *Phys. Rev.* **C72**, 054302 (2005).
31. J. Retamosa, E. Caurier, F. Nowacki and A. Poves, *Phys. Rev.* **C55**, 1266 (1997).
32. G. Audi and A. H. Wapstra, *Nucl. Phys.* **A595**, 409 (1995); R. B. Firestone and C. M. Baglin, *Table of Isotopes*, 8th Edition (John Wiley & Sons, 1998).
33. P. Möller, J. R. Nix and K.-L. Kratz, *LA-UR-94-3898* (1994); *At. Data Nucl. Data Tables* **66**, 131 (1997).
34. B. Blank *et al.*, *Phys. Rev. Lett.* **84**, 1116 (2000).
35. J.A. Clark *et al.*, *Phys. Rev. Lett.* **92**, 192501 (2004).
36. K. D. Sviratcheva, A. I. Georgieva, and J. P. Draayer, *Phys. Rev.* **C69**, 024313 (2004).
37. E. Ormand, *Phys. Rev.* **C55**, 2407 (1997).
38. B. A. Brown, R. R. C. Clement, H. Schatz, and A. Volya, *Phys. Rev.* **C65**, 045802 (2002).
39. M. Honma, T. Mizusaki, and T. Otsuka, *Phys. Rev. Lett.* **77**, 3315 (1996).
40. E. Caurier, F. Novacki, A. Poves, and J. Retamosa, *Phys. Rev. Lett.* **77**, 1954 (1996).
41. P. Van Isacker, O. Juillet, and F. Nowacki, *Phys. Rev. Lett.* **82**, 2060 (1999).
42. K. Kaneko, M. Hasegawa, and T. Mizusaki, *Phys. Rev.* **C66**, 051306(R) (2002).
43. J. P. Vary and D. C. Zheng, “*The Many-Fermion-Dynamics Shell-Model Code*”, Iowa State University, 1994 (unpublished).